

q -Analogues of Convolutions of Fibonacci Numbers

Janine LoBue Tiefenbruck and Jeffrey Remmel

University of California, San Diego

July 8, 2014

The pattern μ

- Avgustinovich, Kitaev, Valyuzhenich: avoidance in permutations
- Jones, Kitaev, Remmel: distribution in cycles of permutations
- LoBue, Remmel: distribution in words



The pattern μ

- Avgustinovich, Kitaev, Valyuzhenich: avoidance in permutations
- Jones, Kitaev, Remmel: distribution in cycles of permutations
- LoBue, Remmel: distribution in words

$$w = w_1 w_2 \dots w_n \in NR([k]), w_i \neq w_{i+1}$$

Definition

The pair $\langle w_i, w_j \rangle$ is a μ -match in w if $i < j$, $w_i < w_j$, and there is no $i < l < j$ such that $w_i \leq w_l \leq w_j$.





The pattern μ

- Avgustinovich, Kitaev, Valyuzhenich: avoidance in permutations
- Jones, Kitaev, Remmel: distribution in cycles of permutations
- LoBue, Remmel: distribution in words

$$w = w_1 w_2 \dots w_n \in NR([k]), w_i \neq w_{i+1}$$

Definition

The pair $\langle w_i, w_j \rangle$ is a μ -match in w if $i < j$, $w_i < w_j$, and there is no $i < l < j$ such that $w_i \leq w_l \leq w_j$.





The pattern μ

- Avgustinovich, Kitaev, Valyuzhenich: avoidance in permutations
- Jones, Kitaev, Remmel: distribution in cycles of permutations
- LoBue, Remmel: distribution in words

$$w = w_1 w_2 \dots w_n \in NR([k]), w_i \neq w_{i+1}$$

Definition

The pair $\langle w_i, w_j \rangle$ is a μ -match in w if $i < j$, $w_i < w_j$, and there is no $i < l < j$ such that $w_i \leq w_l \leq w_j$.



$$N^{i,j}(p, q, t) = \sum_{n \geq 1} N_n^{i,j}(p, q) t^n, \text{ where}$$

$$N_n^{i,j}(p, q) = \sum_{\substack{w \in NR([3]) \\ |w|=n \\ w_1=i, w_n=j}} p^{\text{nontriv}(w)} q^{\text{triv}(w)}$$

A refinement of the generating function for all words in $NR([3])$:

$$N(p, q, t) = 1 + \sum_{i,j \in [3]} N^{i,j}(p, q, t)$$

$$N^{i,j}(p, q, t) = \sum_{n \geq 1} N_n^{i,j}(p, q) t^n, \text{ where}$$

$$N_n^{i,j}(p, q) = \sum_{\substack{w \in NR([3]) \\ |w|=n \\ w_1=i, w_n=j}} p^{\text{nontriv}(w)} q^{\text{triv}(w)}$$

A refinement of the generating function for all words in $NR([3])$:

$$N(p, q, t) = 1 + \sum_{i,j \in [3]} N^{i,j}(p, q, t)$$

Special simplification in the case $k = 3$

The only possible matches are $\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle$. All nontrivial matches are occurrences of the consecutive pattern 13.

- Each word $w \in NR([3])$ belongs to a state of a finite automaton.
- The state records which μ matches may occur when a letter is appended to w .


Example: $w = 2$

$\langle 1, 2 \rangle, \langle 1, 3 \rangle$ matches are not possible

$\langle 2, 3 \rangle$ match is possible

Also in this state is the word $w = 12$.


- Transition from state i to state j if you can append a letter to a word in state i to obtain a word in state j .
- Weight the edges with the change in length (t) and any μ -matches completed by appending the last letter (p 's and q 's).




This gives rise to a system of equations, one for each state.

$$S_3 = t + qtS_2$$

Solve the system for all five states and sum them to get $N(p, q, t)$.



This system of equations gives only the words starting with 3.



This system of equations gives only the words starting with 3.

Important property

All words in a given state have the same last letter.

To get words ending in 1, only sum the shaded states.

$$N^{3,1}(p, q, t) = \frac{t^2 + t^3 - p(1 - q)^2 t^5}{1 - 2qt^2 - q^2 t^3 - pt^2(1 + q^2 t + 2q(q - 1)t^2)}$$

$$N^{3,1}(p, q, t) = \frac{t^2 + t^3 - p(1 - q)^2 t^5}{1 - 2qt^2 - q^2 t^3 - pt^2(1 + q^2 t + 2q(q - 1)t^2)}$$

Theorem

For $n \geq 2$, $N_n^{3,1}(0, 1) = F_{n-2}$.

Proof.

$$\begin{aligned} N^{3,1}(0, 1, t) &= \frac{t^2(1 + t)}{1 - 2qt^2 - q^2 t^3} \\ &= \frac{t^2}{1 - t - t^2} \end{aligned}$$



Theorem

For $n \geq 2$, $N_n^{3,1}(0,1) = F_{n-2}$.

Proof.

Let $\mathcal{C}_n = \{w \in NR([3]) : |w| = n, w_1 = 3, w_n = 1, w \text{ has no } 13s\}$.
Then $N_n^{3,1}(0,1) = |\mathcal{C}_n|$.

$$\mathcal{C}_2 = \{31\} \rightarrow F_0$$

$$\mathcal{C}_3 = \{321\} \rightarrow F_1$$

For $n \geq 4$,

$$\left. \begin{array}{l} 3 \dots 121 \rightarrow \mathbf{3} \dots \mathbf{1}21 \\ 3 \dots 321 \rightarrow \mathbf{3} \dots \mathbf{3}2\mathbf{1} \\ 3 \dots 231 \rightarrow \mathbf{3} \dots \mathbf{2}3\mathbf{1} \end{array} \right\} \rightarrow F_{n-4}$$

$$\left. \begin{array}{l} 3 \dots 321 \rightarrow \mathbf{3} \dots \mathbf{3}2\mathbf{1} \\ 3 \dots 231 \rightarrow \mathbf{3} \dots \mathbf{2}3\mathbf{1} \end{array} \right\} \rightarrow F_{n-3}$$



Theorem

For all $r \geq 0$,

$$N^{3,1}(p, 1, t)|_{p^r} = \frac{t^{2r+2}}{(1-t-t^2)^{r+1}}.$$

so that for all $n, r \geq 0$, $N_n^{3,1}(p, 1)|_{p^r} = \sum_{\substack{j_1, \dots, j_{k+1} \geq 0 \\ n-2r-2 = \sum_{i=1}^{r+1} j_i}} F_{j_1} \cdots F_{j_{r+1}}.$

Proof.

Unique factorization of words with r 13s produces $r+1$ words with no 13s:

$$\boxed{3 \dots 1} \boxed{3 \dots 1} \boxed{3 \dots 1} \boxed{3 \dots 1} \boxed{3 \dots 1}$$

$$N^{3,1}(p, 1, t)|_{p^r} = \prod_{i=1}^{r+1} N^{3,1}(0, 1, t) = \frac{t^{2r+2}}{(1-t-t^2)^{r+1}}.$$



It follows that

① $N_n^{3,1}(0, q)$ is a q -analogue of F_{n-2} , and

② $N_n^{3,1}(p, q)|_{p^r}$ is a q -analogue of $\sum_{\substack{j_1, \dots, j_{r+1} \geq 0 \\ n-2r-2 = \sum_{i=1}^{r+1} j_i}} F_{j_1} \cdots F_{j_{r+1}}.$

n	$N_n^{3,1}(0, q)$	n	$N_n^{3,1}(0, q)$
0	0	8	$12q^3 + q^4$
1	0	9	$8q^3 + 13q^4$
2	1	10	$28q^4 + 6q^5$
3	1	11	$16q^4 + 38q^5 + q^6$
4	$2q$	12	$64q^5 + 25q^6$
5	$2q + q^2$	13	$32q^5 + 104q^6 + 8q^7$
6	$5q^2$	14	$144q^6 + 88q^7 + q^8$
7	$4q^2 + 4q^3$	15	$64q^6 + 272q^7 + 41q^8$

$$N^{3,1}(0, q, t) = \frac{t^2 + t^3}{1 - 2qt^2 - q^2t^3}$$

The denominator gives a recursion that we can also prove combinatorially.

Theorem

For $n \geq 4$, $N_n^{3,1}(0, q) = 2qN_{n-2}^{3,1}(0, q) + q^2N_{n-3}^{3,1}(0, q)$

Proof.

$$\begin{aligned} 3 \dots 121 &\rightarrow \mathbf{3} \dots \mathbf{1}21 \} \rightarrow qN_{n-2}^{3,1} \\ 3 \dots 2321 &\rightarrow \mathbf{3} \dots \mathbf{2}3\mathbf{21} \} \\ 3 \dots 3231 &\rightarrow \mathbf{3} \dots \mathbf{3}2\mathbf{31} \} \rightarrow qN_{n-2}^{3,1} \\ 3 \dots 1231 &\rightarrow \mathbf{3} \dots \mathbf{1}231 \} \rightarrow q^2N_{n-3}^{3,1} \end{aligned}$$



① Patterns in the Coefficients

$$p^{\text{nontriv}(w)} q^{\text{triv}(w)} t^{|w|}$$

n	$N_n^{3,1}(0, q)$	n	$N_n^{3,1}(0, q)$
0	0	8	$12q^3 + q^4$
1	0	9	$8q^3 + 13q^4$
2	1	10	$28q^4 + 6q^5$
3	1	11	$16q^4 + 38q^5 + q^6$
4	$2q$	12	$64q^5 + 25q^6$
5	$2q + q^2$	13	$32q^5 + 104q^6 + 8q^7$
6	$5q^2$	14	$144q^6 + 88q^7 + q^8$
7	$4q^2 + 4q^3$	15	$64q^6 + 272q^7 + 41q^8$

With these initial values and the recursion, we can show

$$N_{2n+1}^{3,1}(0, q) = 2^{n-1} q^{n-1} + \text{higher order terms.}$$

① Patterns in the Coefficients

$$p^{\text{nontriv}(w)} q^{\text{triv}(w)} t^{|w|}$$

n	$N_n^{3,1}(0, q)$	n	$N_n^{3,1}(0, q)$
0	0	8	$12q^3 + q^4$
1	0	9	$8q^3 + 13q^4$
2	1	10	$28q^4 + 6q^5$
3	1	11	$16q^4 + 38q^5 + q^6$
4	$2q$	12	$64q^5 + 25q^6$
5	$2q + q^2$	13	$32q^5 + 104q^6 + 8q^7$
6	$5q^2$	14	$144q^6 + 88q^7 + q^8$
7	$4q^2 + 4q^3$	15	$64q^6 + 272q^7 + 41q^8$

With these initial values and the recursion, we can show

$$N_{2n}^{3,1}(0, q) = (n+2)2^{n-3}q^{n-1} + \text{higher order terms.}$$

① Patterns in the Coefficients

$$p^{\text{nontriv}(w)} q^{\text{triv}(w)} t^{|w|}$$

n	$N_n^{3,1}(0, q)$	n	$N_n^{3,1}(0, q)$
0	0	8	$12q^3 + q^4$
1	0	9	$8q^3 + 13q^4$
2	1	10	$28q^4 + 6q^5$
3	1	11	$16q^4 + 38q^5 + q^6$
4	$2q$	12	$64q^5 + 25q^6$
5	$2q + q^2$	13	$32q^5 + 104q^6 + 8q^7$
6	$5q^2$	14	$144q^6 + 88q^7 + q^8$
7	$4q^2 + 4q^3$	15	$64q^6 + 272q^7 + 41q^8$

$$N_{3n}^{3,1}(0, q) = (1 + 2n(n - 1))q^{2n-2} + \text{lower order terms}$$

$$N_{3n+1}^{3,1}(0, q) = (2n)q^{2n-1} + \text{lower order terms}$$

$$N_{3n+2}^{3,1}(0, q) = q^{2n} + \text{lower order terms}$$

① Patterns in the Coefficients

$$p^{\text{nontriv}(w)} q^{\text{triv}(w)} t^{|w|}$$

n	$N_n^{3,1}(0, q)$	n	$N_n^{3,1}(0, q)$
0	0	8	$12q^3 + q^4$
1	0	9	$8q^3 + 13q^4$
2	1	10	$28q^4 + 6q^5$
3	1	11	$16q^4 + 38q^5 + q^6$
4	$2q$	12	$64q^5 + 25q^6$
5	$2q + q^2$	13	$32q^5 + 104q^6 + 8q^7$
6	$5q^2$	14	$144q^6 + 88q^7 + q^8$
7	$4q^2 + 4q^3$	15	$64q^6 + 272q^7 + 41q^8$

$$N_{2n+3}^{3,1}(0, q)|_{q^{n+k}} = \begin{cases} 0 & \text{if } n < 3k - 2, \\ 1 & \text{if } n = 3k - 2, \\ 4k & \text{if } n = 3k - 1, \\ 2^j \left(4 \binom{2k+j+1}{2k-1} + \binom{2k+j}{2k} \right) & \text{if } n = 3k + j, \quad j \geq 0. \end{cases}$$

$$\begin{aligned}
 N^{3,1}(p, q, t) &= \frac{t^2 + t^3 - p(1 - q)^2 t^5}{1 - 2qt^2 - q^2 t^3 - pt^2(1 + q^2 t + 2q(q - 1)t^2)} \\
 &= \frac{t^2 + t^3 - p(1 - q)^2 t^5}{1 - 2qt^2 - q^2 t^3} \frac{1}{1 - p \left(\frac{t^2(1 + q^2 t + 2q(q - 1)t^2)}{1 - 2qt^2 - q^2 t^3} \right)} \\
 &= \frac{t^2(1 + t)}{1 - 2qt^2 - q^2 t^3} + \\
 &\quad \sum_{r \geq 1} p^r \left(\frac{t^{2r+2}(1 + tq + q(q - 1)t^2)^2(1 + q^2 t + 2q(q - 1)t^2)^{r-1}}{(1 - 2qt^2 - q^2 t^3)^{r+1}} \right)
 \end{aligned}$$

② $N_n^{3,1}(p, q)|_{p^r}$ when $r = 1$

$p^{nontriv(w)} q^{triv(w)} t^{|w|}$

$$N^{3,1}(p, q, t)|_p = \frac{t^4(1 + tq + q(q-1)t^2)^2}{(1 - 2qt^2 - q^2t^3)^2}$$

In this case, $N_n^{3,1}(p, q)|_p$ is a q -analogue of $\sum_{j=0}^{n-4} F_j F_{n-4-j}$.

n	$N_n^{3,1}(p, q) _p$	n	$N_n^{3,1}(p, q) _p$
0	0	9	$24q^3 + 14q^4$
1	0	10	$12q^3 + 51q^4 + 8q^5$
2	0	11	$66q^4 + 62q^5 + 2q^6$
3	0	12	$28q^4 + 162q^5 + 45q^6$
4	1	13	$172q^5 + 230q^6 + 18q^7$
5	$2q$	14	$64q^5 + 475q^6 + 202q^7 + 3q^8$
6	$2q + 3q^2$	15	$432q^6 + 768q^7 + 108q^8$
7	$8q^2 + 2q^3$	16	$144q^6 + 1320q^7 + 789q^8 + 32q^9$
8	$5q^2 + 14q^3 + q^4$	17	$1056q^7 + 2388q^7 + 522q^9 + 4q^{10}$

② Patterns in the Coefficients

$$p^{nontriv(w)} q^{triv(w)} t^{|w|}$$

n	$N_n^{3,1}(p, q) _p$	n	$N_n^{3,1}(p, q) _p$
0	0	9	$24q^3 + 14q^4$
1	0	10	$12q^3 + 51q^4 + 8q^5$
2	0	11	$66q^4 + 62q^5 + 2q^6$
3	0	12	$28q^4 + 162q^5 + 45q^6$
4	1	13	$172q^5 + 230q^6 + 18q^7$
5	$2q$	14	$64q^5 + 475q^6 + 202q^7 + 3q^8$
6	$2q + 3q^2$	15	$432q^6 + 768q^7 + 108q^8$
7	$8q^2 + 2q^3$	16	$144q^6 + 1320q^7 + 789q^8 + 32q^9$
8	$5q^2 + 14q^3 + q^4$	17	$1056q^7 + 2388q^7 + 522q^9 + 4q^{10}$

$$N_{2n+4}^{3,1}(p, q)|_p = (n+3)2^{n-2}q^n + \text{higher order terms}$$

$$N_{2n+5}^{3,1}(p, q)|_p = (18 + 13n + n^2)2^{n-3}q^{n+1} + \text{higher order terms}$$

② Patterns in the Coefficients

$$p^{nontriv(w)} q^{triv(w)} t^{|w|}$$

n	$N_n^{3,1}(p, q) _p$	n	$N_n^{3,1}(p, q) _p$
0	0	9	$24q^3 + 14q^4$
1	0	10	$12q^3 + 51q^4 + 8q^5$
2	0	11	$66q^4 + 62q^5 + 2q^6$
3	0	12	$28q^4 + 162q^5 + 45q^6$
4	1	13	$172q^5 + 230q^6 + 18q^7$
5	$2q$	14	$64q^5 + 475q^6 + 202q^7 + 3q^8$
6	$2q + 3q^2$	15	$432q^6 + 768q^7 + 108q^8$
7	$8q^2 + 2q^3$	16	$144q^6 + 1320q^7 + 789q^8 + 32q^9$
8	$5q^2 + 14q^3 + q^4$	17	$1056q^7 + 2388q^7 + 522q^9 + 4q^{10}$

$$N_{3n+3}^{3,1}(p, q)|_p = n(2n^2 - 2n + 3)q^{2n} + \text{lower order terms}$$

$$N_{3n+4}^{3,1}(p, q)|_p = 2n^2q^{2n+1} + \text{lower order terms}$$

$$N_{3n+5}^{3,1}(p, q)|_p = nq^{2n+2} + \text{lower order terms}$$

Theorem

$$N_n^{1,1}(0, 1) = N_n^{3,3}(0, 1) = F_{n-3}$$

$$N_n^{2,1}(0, 1) = N_n^{3,2}(0, 1) = F_{n-2}$$

$$N_n^{1,2}(0, 1) = N_n^{2,3}(0, 1) = F_{n-3} + (-1)^n$$

$$N_n^{1,3}(0, 1) = F_{n-4} + (-1)^{n-1}$$

$$N_n^{2,2}(0, 1) = F_{n-2} + (-1)^{n-1}$$

$$N^{1,1}(0, q, t) = \frac{t - 2qt^3 - q^2t^4}{1 - 3qt^2 - 2q^2t^3 - (q^3 - q^2)t^4}$$

n	$N_n^{1,1}(0, q)$	$N_n^{1,1}(0, 1)$	n	$N_n^{1,1}(0, q)$	$N_n^{1,1}(0, 1)$
0	0	0	6	$5q^3$	5
1	1	1	7	$5q^3 + 6q^4$	11
2	0	0	8	$18q^4 + 3q^5$	21
3	q	1	9	$13q^4 + 29q^5 + q^6$	43
4	q^2	1	10	$59q^5 + 26q^6$	85
5	$2q^2 + q^3$	3	11	$34q^5 + 122q^6 + 15q^7$	171

Jacobsthal numbers: $J_n = J_{n-1} + 2J_{n-2}$

$$k = 4$$

$$p^{\text{nontriv}(w)} q^{\text{triv}(w)} t^{|w|}$$

$$N^{1,1}(0, q, t) = \frac{t - 2qt^3 - q^2t^4}{1 - 3qt^2 - 2q^2t^3 - (q^3 - q^2)t^4}$$

n	$N_n^{1,1}(0, q)$	$N_n^{1,1}(0, 1)$	n	$N_n^{1,1}(0, q)$	$N_n^{1,1}(0, 1)$
0	0	0	6	$5q^3$	5
1	1	1	7	$5q^3 + 6q^4$	11
2	0	0	8	$18q^4 + 3q^5$	21
3	q	1	9	$13q^4 + 29q^5 + q^6$	43
4	q^2	1	10	$59q^5 + 26q^6$	85
5	$2q^2 + q^3$	3	11	$34q^5 + 122q^6 + 15q^7$	171

Jacobsthal numbers: $J_n = J_{n-1} + 2J_{n-2}$

$$N_{2n+1}^{1,1}(0, q) = F_{2n+1}q^n + \text{higher order terms}$$

