

Functional equations for enumerating 1324-avoiding permutations

Brian Nakamura

Rutgers University

Permutation Patterns 2014

Upcoming...

Upcoming...

DIMACS Conference on Challenges of Identifying Integer Sequences
Celebrating 50 years of the OEIS and Neil Sloane's 75th birthday!
October 9–10, 2014

Poster abstracts are welcomed!

Some funding may be available for “early career” researchers to attend.

For more details, see: <http://dimacs.rutgers.edu/Workshops/OEIS/>

Pattern occurrences

Definition

Given a (permutation) pattern $\tau = \tau_1 \dots \tau_k$, we say that permutation $\pi = \pi_1 \dots \pi_n$ *contains* the pattern τ if there exists $1 \leq i_1 < \dots < i_k \leq n$ such that $\pi_{i_1} \pi_{i_2} \dots \pi_{i_k}$ is order-isomorphic to τ .

Example

If pattern $\tau = 123$,

- $\pi = 54321$ has zero occurrences of τ ,
- $\pi = 42135$ has two occurrences of τ .

Some notation

Definition

Given a pattern $\tau \in \mathcal{S}_k$, define

$$\begin{aligned}\mathcal{S}_n(\tau) &:= \{\pi \in \mathcal{S}_n : \pi \text{ avoids } \tau\}, \\ s_n(\tau) &:= |\mathcal{S}_n(\tau)|.\end{aligned}$$

Some notation

Definition

Given a pattern $\tau \in \mathcal{S}_k$, define

$$\begin{aligned}\mathcal{S}_n(\tau) &:= \{\pi \in \mathcal{S}_n : \pi \text{ avoids } \tau\}, \\ s_n(\tau) &:= |\mathcal{S}_n(\tau)|.\end{aligned}$$

Definition

Given a pattern τ and $r \geq 0$, define

$$\begin{aligned}\mathcal{S}_n(\tau, r) &:= \{\pi \in \mathcal{S}_n : \pi \text{ with exactly } r \text{ occurrences of } \tau\}, \\ s_n(\tau, r) &:= |\mathcal{S}_n(\tau, r)|.\end{aligned}$$

Pattern avoiding permutations ($r = 0$)

The $r = 0$ case (classical pattern avoidance) has received a lot of attention.

This gets difficult very quickly.

Even for length 4 patterns, many unresolved questions remain (e.g., pattern 1324).

Regarding 1324-avoiders...

Regarding 1324-avoiders...

“Conjecture” (Zeilberger, 2005)

“Not even God knows $s_{1000}(1324)$.”

Regarding 1324-avoiders...

“Conjecture” (Zeilberger, 2005)

“Not even God knows $s_{1000}(1324)$.”

“Conjecture” (Steingrímsson, 2013)

“I’m not sure how good Zeilberger’s God is at math, but I believe that some humans will find this number in the not so distant future.”

Regarding 1324-avoiders...

“Conjecture” (Zeilberger, 2005)

“Not even God knows $s_{1000}(1324)$.”

“Conjecture” (Steingrímsson, 2013)

“I’m not sure how good Zeilberger’s God is at math, but I believe that some humans will find this number in the not so distant future.”

FYI: There’s now a bet whether this will be found by year 2030.

Exactly r copies of a pattern ($r > 0$)

Most work on $r > 0$ focuses on length 3 patterns:

- $s_n(132, r)$ studied by Bóna, Mansour and Vainshtein, Fulmek, and others.
- $s_n(123, r)$ studied by Noonan and Zeilberger, Fulmek, Callan, and others.

Exactly r copies of a pattern ($r > 0$)

Most work on $r > 0$ focuses on length 3 patterns:

- $s_n(132, r)$ studied by Bóna, Mansour and Vainshtein, Fulmek, and others.
- $s_n(123, r)$ studied by Noonan and Zeilberger, Fulmek, Callan, and others.

GOAL: for fixed pattern τ and fixed r , compute $s_n(\tau, r)$ “quickly”.

Exactly r copies of a pattern ($r > 0$)

Most work on $r > 0$ focuses on length 3 patterns:

- $s_n(132, r)$ studied by Bóna, Mansour and Vainshtein, Fulmek, and others.
- $s_n(123, r)$ studied by Noonan and Zeilberger, Fulmek, Callan, and others.

GOAL: for fixed pattern τ and fixed r , compute $s_n(\tau, r)$ “quickly”.

For motivation, we will start with $\tau = 12 \dots k$.

Additional definitions

Assume pattern $\tau = 123$ (equiv. abc).

Additional definitions

Assume pattern $\tau = 123$ (equiv. abc).

Definition

For variables $t, x_1, \dots, x_n$, define

$$\text{weight}(\pi) := t^{\# \text{ of } abc \text{ in } \pi} \cdot \prod_{i=1}^n x_i^{\# \text{ of } ab \text{ in } \pi \text{ s.t. } a=i}$$

$$P_n(t; x_1, \dots, x_n) := \sum_{\pi \in \mathcal{S}_n} \text{weight}(\pi)$$

Additional definitions

Assume pattern $\tau = 123$ (equiv. abc).

Definition

For variables $t, x_1, \dots, x_n$, define

$$\text{weight}(\pi) := t^{\# \text{ of } abc \text{ in } \pi} \cdot \prod_{i=1}^n x_i^{\# \text{ of } ab \text{ in } \pi \text{ s.t. } a=i}$$

$$P_n(t; x_1, \dots, x_n) := \sum_{\pi \in \mathcal{S}_n} \text{weight}(\pi)$$

Observe: coeff. of t^r in $P_n(t; 1, \dots, 1) = s_n(123, r)$.

Functional equations for 123

Theorem (Noonan and Zeilberger, 1996)

$$P_n(t; x_1, \dots, x_n) = \sum_{i=1}^n x_i^{n-i} P_{n-1}(t; x_1, \dots, x_{i-1}, tx_{i+1}, \dots, tx_n)$$

Functional equations for 123

Theorem (Noonan and Zeilberger, 1996)

$$P_n(t; x_1, \dots, x_n) = \sum_{i=1}^n x_i^{n-i} P_{n-1}(t; x_1, \dots, x_{i-1}, tx_{i+1}, \dots, tx_n)$$

We can use this functional equation to compute $P_n(t; 1, \dots, 1)$.

NOTE: can apply this directly to $P_n(t; 1, \dots, 1)$

Functional equations for 123

Theorem (Noonan and Zeilberger, 1996)

$$P_n(t; x_1, \dots, x_n) = \sum_{i=1}^n x_i^{n-i} P_{n-1}(t; x_1, \dots, x_{i-1}, tx_{i+1}, \dots, tx_n)$$

We can use this functional equation to compute $P_n(t; 1, \dots, 1)$.

NOTE: can apply this directly to $P_n(t; 1, \dots, 1)$

For example, if $x_1 = \dots = x_n = 1$, then

$$P_n(t; 1 [n \text{ times}]) = \sum_{i=1}^n P_{n-1}(t; 1 [i-1 \text{ times}], t [n-i \text{ times}])$$

Maple implementation

Can apply other computational methods to quickly find coeff. of t^r in $P_n(t; 1, \dots, 1)$.

Maple implementation

Can apply other computational methods to quickly find coeff. of t^r in $P_n(t; 1, \dots, 1)$.

For fixed r , it can be implemented to run in polynomial-time (but memory-intensive!).

Maple implementation

Can apply other computational methods to quickly find coeff. of t^r in $P_n(t; 1, \dots, 1)$.

For fixed r , it can be implemented to run in polynomial-time (but memory-intensive!).

Example

- For $r = 0$, the first 10 terms of $s_n(123, r)$ are:
1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796
- For $r = 1$, the first 10 terms of $s_n(123, r)$ are:
0, 0, 1, 6, 27, 110, 429, 1638, 6188, 23256
- For $r = 6$, the values of $s_n(123, r)$ for $15 \leq n \leq 20$ are:
327200581, 1501719377, 6773007550, 30100185693,
132099138291, 573518305776

Extending the approach to $\tau = 1234$

Definition

For variables $t, x_1, \dots, x_n, y_1, \dots, y_n$, define

$$\text{weight}(\pi) := t^{\# \text{ of } abcd \text{ in } \pi} \cdot \prod_{i=1}^n x_i^{\# \text{ of } ab \text{ in } \pi \text{ s.t. } a=i} \cdot \prod_{j=1}^n y_j^{\# \text{ of } abc \text{ in } \pi \text{ s.t. } a=j}$$

$$P_n(t; x_1, \dots, x_n; y_1, \dots, y_n) := \sum_{\pi \in \mathcal{S}_n} \text{weight}(\pi)$$

Observe: coeff. of t^r in $P_n(t; 1, \dots, 1; 1, \dots, 1) = s_n(1234, r)$.

Functional equations for 1234

Theorem (Noonan and Zeilberger, 1996)

$$P_n(t; x_1, \dots, x_n; y_1, \dots, y_n) = \sum_{i=1}^n x_i^{n-i} P_{n-1}(t; x_1, \dots, x_{i-1}, y_i x_{i+1}, \dots, y_i x_n; y_1, \dots, y_{i-1}, t y_{i+1}, \dots, t y_n)$$

We can use this functional equation to compute $P_n(t; 1, \dots, 1; 1, \dots, 1)$.

Computing some values

Similar computational methods can be applied to this case.

Computing some values

Similar computational methods can be applied to this case.

Again, this was implemented in Maple.

Computing some values

Similar computational methods can be applied to this case.

Again, this was implemented in Maple.

Example

- We can compute $s_n(1234, 1)$ up to $n = 70$:
 $s_{70}(1234, 1) \approx 4.335 \times 10^{59}$ (could not guess a recurrence...)

Computing some values

Similar computational methods can be applied to this case.

Again, this was implemented in Maple.

Example

- We can compute $s_n(1234, 1)$ up to $n = 70$:
 $s_{70}(1234, 1) \approx 4.335 \times 10^{59}$ (could not guess a recurrence...)
- With help from Manuel Kauers, we computed $s_n(1234, 1)$ up to $n = 200$:
 $s_{200}(1234, 1) \approx 6.595 \times 10^{181}$ (but still no recurrence...)

Computing some values

Similar computational methods can be applied to this case.

Again, this was implemented in Maple.

Example

- We can compute $s_n(1234, 1)$ up to $n = 70$:
 $s_{70}(1234, 1) \approx 4.335 \times 10^{59}$ (could not guess a recurrence...)
- With help from Manuel Kauers, we computed $s_n(1234, 1)$ up to $n = 200$:
 $s_{200}(1234, 1) \approx 6.595 \times 10^{181}$ (but still no recurrence...)

This approach extendable to any increasing pattern $12 \dots k$.

Further extensions

Previous approach is extendable to 132, 1243, and more generally $12 \dots (k-2)(k)(k-1)$.

Further extensions

Previous approach is extendable to 132, 1243, and more generally $12 \dots (k-2)(k)(k-1)$.

This approach can be generalized to handle other patterns by considering more complicated catalytic variables $x_{i,j}$'s.

Further extensions

Previous approach is extendable to 132, 1243, and more generally $12 \dots (k-2)(k)(k-1)$.

This approach can be generalized to handle other patterns by considering more complicated catalytic variables $x_{i,j}$'s.

Some additional patterns that can be handled with this approach:

- Patterns 231, 2341, and more generally $23 \dots k1$
(Recall Vatter's talk and the comment on *adcb*)

Further extensions

Previous approach is extendable to 132, 1243, and more generally $12 \dots (k-2)(k)(k-1)$.

This approach can be generalized to handle other patterns by considering more complicated catalytic variables $x_{i,j}$'s.

Some additional patterns that can be handled with this approach:

- Patterns 231, 2341, and more generally $23 \dots k1$
(Recall Vatter's talk and the comment on *adcb*)
- Pattern 1324

Set-up for 1324

We consider the catalytic variables:

- $x_{i,j}$ ($1 \leq i \leq j \leq n$)
- $y_{i,j}$ ($1 \leq j \leq i \leq n$).

Set-up for 1324

We consider the catalytic variables:

- $x_{i,j}$ ($1 \leq i \leq j \leq n$)
- $y_{i,j}$ ($1 \leq j \leq i \leq n$).

Variables $x_{i,j}$ will be written as a matrix of variables:

$$X_n := \begin{bmatrix} x_{1,1} & \cdots & x_{1,n} \\ & \ddots & \\ \vdots & x_{i,i} & \vdots \\ & & \ddots & \\ x_{n,1} & \cdots & x_{n,n} \end{bmatrix}$$

(similarly for variables $y_{i,j}$ and matrix Y_n)

Weight function for 1324

Definition

For variables t , $x_{i,j}$'s, and $y_{i,j}$'s ($1 \leq i, j \leq n$), define

$$\text{weight}(\pi) := t^{\# \text{ of } acbd \text{ in } \pi} \cdot \prod_{i \leq j} x_{i,j}^{\# \text{ of } ab \text{ in } \pi \text{ s.t. } a=i, b>j} \cdot \prod_{j \leq i} y_{i,j}^{\# \text{ of } bac \text{ in } \pi \text{ s.t. } b=i, a \geq j}$$

Weight function for 1324

Definition

For variables t , $x_{i,j}$'s, and $y_{i,j}$'s ($1 \leq i, j \leq n$), define

$$\text{weight}(\pi) := t^{\# \text{ of } acbd \text{ in } \pi} \cdot \prod_{i \leq j} x_{i,j}^{\# \text{ of } ab \text{ in } \pi \text{ s.t. } a=i, b>j} \cdot \prod_{j \leq i} y_{i,j}^{\# \text{ of } bac \text{ in } \pi \text{ s.t. } b=i, a \geq j}$$

Definition

$$P_n(t; X_n; Y_n) := \sum_{\pi \in \mathcal{S}_n} \text{weight}(\pi)$$

Weight function for 1324

Definition

For variables t , $x_{i,j}$'s, and $y_{i,j}$'s ($1 \leq i, j \leq n$), define

$$\text{weight}(\pi) := t^{\# \text{ of } acbd \text{ in } \pi} \cdot \prod_{i \leq j} x_{i,j}^{\# \text{ of } ab \text{ in } \pi \text{ s.t. } a=i, b>j} \cdot \prod_{j \leq i} y_{i,j}^{\# \text{ of } bac \text{ in } \pi \text{ s.t. } b=i, a \geq j}$$

Definition

$$P_n(t; X_n; Y_n) := \sum_{\pi \in \mathcal{S}_n} \text{weight}(\pi)$$

Observe: coeff. of t^r in $P_n(t; \mathbf{1}_n; \mathbf{1}_n) = s_n(1324, r)$.

A matrix operator

Define operator R_1 on an arbitrary $n \times n$ square matrix Y_n and $1 \leq i < n$:

$$R_1(Y_n, i) := \begin{bmatrix} y_{1,1} & \cdots & y_{1,i-1} & u_1 & y_{1,i+2} & \cdots & y_{1,n} \\ \vdots & \ddots & & \vdots & & & \vdots \\ y_{i-1,1} & \cdots & y_{i-1,i-1} & u_{i-1} & y_{i-1,i+2} & \cdots & y_{i-1,n} \\ y_{i+1,1} & \cdots & y_{i+1,i-1} & u_{i+1} & y_{i+1,i+2} & \cdots & y_{i+1,n} \\ \vdots & & & \vdots & & \ddots & \vdots \\ y_{n,1} & \cdots & y_{n,i-1} & u_n & y_{n,i+2} & \cdots & y_{n,n} \end{bmatrix}$$

with

$$u_k := t \cdot y_{k,i} \cdot y_{k,i+1}$$

And $R_1(Y_n, n)$ defined to be Y_n with n -th row and n -th column deleted.

... and another

Define operator R_2 on two $n \times n$ square matrices X_n and Y_n and $1 < i \leq n$:

$$R_2(X_n, Y_n, i) := \begin{bmatrix} x_{1,1} & \cdots & x_{1,i-2} & w_1 & x_{1,i+1} & \cdots & x_{1,n} \\ \vdots & \ddots & & \vdots & & & \vdots \\ x_{i-1,1} & \cdots & x_{i-1,i-2} & w_{i-1} & x_{i-1,i+1} & \cdots & x_{i-1,n} \\ x_{i+1,1} & \cdots & x_{i+1,i-2} & w_{i+1} & x_{i+1,i+1} & \cdots & x_{i+1,n} \\ \vdots & & & \vdots & & \ddots & \vdots \\ x_{n,1} & \cdots & x_{n,i-2} & w_n & x_{n,i+1} & \cdots & x_{n,n} \end{bmatrix}$$

with

$$w_k := y_{i,1}y_{i,2} \cdots y_{i,k} \cdot x_{k,i-1} \cdot x_{k,i}$$

And $R_2(X_n, Y_n, 1)$ is X_n with the 1-st row and 1-st column deleted.

Functional equation for 1324

Theorem

$$P_n(t; X_n, Y_n) = \sum_{i=1}^n x_{i,i}^{n-i} x_{i,i+1}^{n-i-1} \cdots x_{i,n-1}^1 \cdot P_{n-1}(t; R_2(X_n, Y_n, i); R_1(Y_n, i)).$$

And the coeff. of t^r in $P_n(t; \mathbf{1}_n, \mathbf{1}_n)$ is exactly $s_n(1324, r)$.

Avoiding 1324

We can specialize the functional equation for the $r = 0$ case.

Avoiding 1324

We can specialize the functional equation for the $r = 0$ case.

This allows us to compute the first 31 terms. (A061552 in OEIS)
(joint work with F. Johansson)

For example, $s_{31}(1324) = 1914189093351633702834757$.

Avoiding 1324

We can specialize the functional equation for the $r = 0$ case.

This allows us to compute the first 31 terms. (A061552 in OEIS)
(joint work with F. Johansson)

For example, $s_{31}(1324) = 1914189093351633702834757$.

Easy to refine by the number of inversions.

(e.g., confirmed CJS conjecture on 1324-avoiders with exactly k inversions up to n , $k \leq 23$)

Some other recent work

In recent work, Conway and Guttman provide a refinement of the previous algorithm.

Some other recent work

In recent work, Conway and Guttmann provide a refinement of the previous algorithm.

They compute $s_n(1324)$ up to $n = 36$ and use this to empirically estimate asymptotics.

Some other recent work

In recent work, Conway and Guttmann provide a refinement of the previous algorithm.

They compute $s_n(1324)$ up to $n = 36$ and use this to empirically estimate asymptotics.

They conjecture:

$$s_n(1324) \sim B \cdot \mu^n \cdot \mu_1^{\sqrt{n}} \cdot n^g$$

with $B = 9.5 \pm 1.0$, $\mu = 11.60 \pm 0.01$, $\mu_1 = 0.0398 \pm 0.0010$, and $g = -1.1 \pm 0.2$.

Thank you

Thank you!