

A general theory of Wilf-equivalence
for permutation classes $\mathcal{A}_V(231, \pi)$,
and other Catalan structures

Mathilde Bouvel (Universität Zürich)

joint work with Michael Albert (University of Otago)

Permutation Patterns 2014

Wilf-equivalence

- $\text{Av}(B)$ is the class of permutations avoiding all patterns in B .
- (a_n) where $a_n = |\text{Av}_n(B)|$ is its enumeration sequence.
- Its generating function is $\sum a_n t^n$.
- Two permutation classes $\text{Av}(B_1)$ and $\text{Av}(B_2)$ are **Wilf-equivalent** if they have the same enumeration sequence (or generating function).

Wilf-equivalence

- $\text{Av}(B)$ is the class of permutations avoiding all patterns in B .
- (a_n) where $a_n = |\text{Av}_n(B)|$ is its enumeration sequence.
- Its generating function is $\sum a_n t^n$.
- Two permutation classes $\text{Av}(B_1)$ and $\text{Av}(B_2)$ are **Wilf-equivalent** if they have the same enumeration sequence (or generating function).

More or less famous Wilf-equivalences:

- $\text{Av}(123)$ and $\text{Av}(231)$, enumerated by the Catalan numbers Cat_n
- There are three Wilf-equivalence classes for permutation classes $\text{Av}(\pi)$ with π of size 4, the enumeration of $\text{Av}(1324)$ being open.
- Check all Wilf-equivalences between $\text{Av}(\pi, \tau)$ when π and τ have size 3 or 4 on Wikipedia.
- Some results for arbitrary long patterns:
 $\text{Av}(231 \oplus \pi)$ and $\text{Av}(312 \oplus \pi)$ [West & Stankova 02]

Wilf-equivalences of permutation classes $\text{Av}(231, \pi)$

Harmless assumption: In $\text{Av}(231, \pi)$ throughout the talk, π avoids 231.

Wilf-equivalences of permutation classes $\text{Av}(231, \pi)$

Harmless assumption: In $\text{Av}(231, \pi)$ throughout the talk, π avoids 231.

Known results: Three families of patterns π such that the generating function of $\text{Av}(231, \pi)$ is C_n , where $n = |\pi|$, defined by $C_0 = 1$,
 $C_{n+1} = \frac{1}{1-tC_n}$ [Mansour & Vainshtein 01+02; Albert & Bouvel 13]

Remark: The generating functions C_n are truncations at level n of the continued fraction defining the generating function of Catalan numbers:

$$C = \frac{1}{1 - \frac{t}{1 - \frac{t}{1 - \frac{t}{1 - \dots}}}}.$$

Wilf-equivalences of permutation classes $\text{Av}(231, \pi)$

Harmless assumption: In $\text{Av}(231, \pi)$ throughout the talk, π avoids 231.

Known results: Three families of patterns π such that the generating function of $\text{Av}(231, \pi)$ is C_n , where $n = |\pi|$, defined by $C_0 = 1$,
 $C_{n+1} = \frac{1}{1-t} C_n$ [Mansour & Vainshtein 01+02; Albert & Bouvel 13]

New results on this topic:

- Description of all patterns π of size n such that the generating function of $\text{Av}(231, \pi)$ is C_n .
- There are $\text{Motz}_n = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} \text{Cat}_k$ such patterns.
- Bijections between $\text{Av}(231, \pi)$ and $\text{Av}(231, \pi')$ for any such patterns.
- For τ of size n , the generating function of $\text{Av}(231, \tau)$ either is C_n or C_n dominates it term by term (and eventually strictly).

Wilf-equivalences of permutation classes $\text{Av}(231, \pi)$

Harmless assumption: In $\text{Av}(231, \pi)$ throughout the talk, π avoids 231.

Known results: Three families of patterns π such that the generating function of $\text{Av}(231, \pi)$ is C_n , where $n = |\pi|$, defined by $C_0 = 1$,
 $C_{n+1} = \frac{1}{1-t} C_n$ [Mansour & Vainshtein 01+02; Albert & Bouvel 13]

New results on this topic:

- Description of all patterns π of size n such that the generating function of $\text{Av}(231, \pi)$ is C_n .
- There are $\text{Motz}_n = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} \text{Cat}_k$ such patterns.
- Bijections between $\text{Av}(231, \pi)$ and $\text{Av}(231, \pi')$ for any such patterns.
- For τ of size n , the generating function of $\text{Av}(231, \tau)$ either is C_n or C_n dominates it term by term (and eventually strictly).


Main objective: Find all Wilf-equivalences between classes $\text{Av}(231, \pi)$.

Substructures in Catalan objects


Catalan structures, and their substructures

- 231-avoiding permutations


41327658 =




- Plane forests




- Complete binary trees



- Dyck paths




- Arch systems




Catalan structures, and their substructures

- 231-avoiding permutations


41327658 =




- Plane forests



- Complete binary trees



- Dyck paths



- Arch systems




Catalan structures, and their substructures

- 231-avoiding permutations


31254 =



- Dyck paths




- Plane forests



- Arch systems



- Complete binary trees




Catalan structures, and their substructures

- 231-avoiding permutations


31254 =




- Plane forests



- Complete binary trees



- Dyck paths



- Arch systems




Catalan structures, and their substructures

- 231-avoiding permutations


31254 =




- Plane forests



- Complete binary trees



- Dyck paths



- Arch systems




Catalan structures, and their substructures

- 231-avoiding permutations


31254 =



- Dyck paths




- Plane forests



- Arch systems



- Complete binary trees




Catalan structures, and their substructures

- 231-avoiding permutations


31254 =



- Dyck paths




- Plane forests



- Arch systems



- Complete binary trees




Catalan structures, and their substructures

- 231-avoiding permutations


31254 =




- Plane forests



- Complete binary trees



- Dyck paths




- Arch systems



Fact: The usual bijections relating our quartet of Catalan structures preserve the substructure relation.

Classes of arch systems instead of $\text{Av}(231, \pi)$

- The following bijection between 231-avoiding permutations and arch systems preserves the substructure relation:



- Therefore, for all π avoiding 231, $\text{Av}(231, \pi) \xleftrightarrow{\varphi} \text{Av}(\varphi(\pi))$.
- We will study classes $\text{Av}(A)$ of arch systems avoiding some subsystem A , but all results can be translated to classes $\text{Av}(231, \pi)$ via φ .

Questions addressed in this talk

- Which arch systems A are Wilf-equivalent?
i.e. which classes $A_V(A)$ have the same enumeration?
- Bijections between $A_V(A)$ and $A_V(B)$ for Wilf-equivalent arch systems A and B ?
- How many Wilf-equivalence classes of arch systems are there?

Questions addressed in this talk

- Which arch systems A are **Wilf-equivalent**?
i.e. which classes $A_V(A)$ have the same enumeration?
- **Bijections** between $A_V(A)$ and $A_V(B)$ for Wilf-equivalent arch systems A and B ?
- **How many** Wilf-equivalence classes of arch systems are there?

Observation and terminology:

An arch system is a concatenation of **atoms**, i.e. (non-empty) arch systems having a single outermost arch.



**An equivalence relation
strongly related to Wilf-equivalence**

An equivalence relation on arch systems

The binary relation, $\sim$, is the finest equivalence relation that satisfies:

$$(0) \quad A \sim A$$

$$(1) \quad A \sim B \implies \overline{A} \sim \overline{B}$$

$$(2) \quad a \sim b \implies PaQ \sim PbQ$$

$$(3) \quad PabQ \sim PbaQ$$

$$(4) \quad a\overline{bc} \sim \overline{ab}c$$

where A , B , P and Q denote arbitrary arch systems
and a , b and c denote atoms or empty arch systems.

An equivalence relation on arch systems

The binary relation, $\sim$, is the finest equivalence relation that satisfies:

$$(0) \quad A \sim A$$

$$(1) \quad A \sim B \implies \overline{A} \sim \overline{B}$$

$$(2) \quad a \sim b \implies PaQ \sim PbQ$$

$$(3) \quad PabQ \sim PbaQ$$

$$(4) \quad a\overline{bc} \sim \overline{ab}c$$

where A , B , P and Q denote arbitrary arch systems
and a , b and c denote atoms or empty arch systems.

Terminology: The equivalence classes of $\sim$ are called **cohorts**.

$\sim$ is (a refinement of?) Wilf-equivalence

Main theorem: If A and B are arch systems such that $A \sim B$ then $A_V(A)$ and $A_V(B)$ have the same enumeration, *i.e.* are Wilf-equivalent.

$\sim$ is (a refinement of?) Wilf-equivalence

Main theorem: If A and B are arch systems such that $A \sim B$ then $A_V(A)$ and $A_V(B)$ have the same enumeration, *i.e.* are Wilf-equivalent.

Conjecture: $\sim$ coincides with Wilf-equivalence.

Data, obtained with PermLab:

The conjecture holds for arch systems of size up to 15 (where $\sim$ has 16,709 equivalence classes on the $Cat_{15} = 9,694,845$ arch systems).

$\sim$ is (a refinement of?) Wilf-equivalence

Main theorem: If A and B are arch systems such that $A \sim B$ then $\text{Av}(A)$ and $\text{Av}(B)$ have the same enumeration, i.e. are Wilf-equivalent.

Conjecture: $\sim$ coincides with Wilf-equivalence.

Data, obtained with PermLab:

The conjecture holds for arch systems of size up to 15 (where $\sim$ has 16,709 equivalence classes on the $\text{Cat}_{15} = 9,694,845$ arch systems).

Additional results:

- Asymptotic enumeration of the number of cohorts.
- One cohort of arch systems of size n (conjecturally the biggest one) contains Motz_n arch systems, and for A in this cohort $\text{Av}(A)$ is enumerated by C_n .
- Comparison of the enumeration sequences of $\text{Av}(A)$ and $\text{Av}(B)$.

Idea of the proof

Overview of the proof

Main theorem: If A and B are arch systems such that $A \sim B$ then $A_V(A)$ and $A_V(B)$ have the same enumeration, *i.e.* are Wilf-equivalent.

Overview of the proof... by induction!

Main theorem: If A and B are arch systems such that $A \sim B$ then $A_V(A)$ and $A_V(B)$ have the same enumeration, i.e. are Wilf-equivalent.

Base case: If $A = B$ then $A_V(A)$ and $A_V(B)$ are Wilf-equivalent...

Inductive case: One case for each rule defining $\sim$.

Rule	bijjective proof	analytic proof
(1) $A \sim B \implies \overline{A} \sim \overline{B}$	yes	-
(2) $a \sim b \implies PaQ \sim PbQ$	yes	-
(3) $PabQ \sim PbaQ$	yes	-
(4) $a\overline{bc} \sim \overline{ab}c$	no	yes

Overview of the proof... by induction!

Main theorem: If A and B are arch systems such that $A \sim B$ then $A_V(A)$ and $A_V(B)$ have the same enumeration, i.e. are Wilf-equivalent.

Base case: If $A = B$ then $A_V(A)$ and $A_V(B)$ are Wilf-equivalent...

Inductive case: One case for each rule defining $\sim$.

Rule	bijjective proof	analytic proof
(1) $A \sim B \implies \overline{A} \sim \overline{B}$	yes	-
(2) $a \sim b \implies PaQ \sim PbQ$	yes	-
(3) $PabQ \sim PbaQ$	yes	-
(4) $a\overline{bc} \sim \overline{ab}c$	no	yes
(4 weak) $a\overline{b} \sim \overline{ba}$	yes	-

Overview of the proof... by induction!

Main theorem: If A and B are arch systems such that $A \sim B$ then $A_V(A)$ and $A_V(B)$ have the same enumeration, i.e. are Wilf-equivalent.

Base case: If $A = B$ then $A_V(A)$ and $A_V(B)$ are Wilf-equivalent...

Inductive case: One case for each rule defining $\sim$.

Rule	bijective proof	analytic proof
(1) $A \sim B \implies \overline{A} \sim \overline{B}$	yes	-
(2) $a \sim b \implies PaQ \sim PbQ$	yes	-
(3) $PabQ \sim PbaQ$	yes	-
(4) $a\overline{bc} \sim \overline{ab}c$	no	yes
(4 weak) $a\overline{b} \sim \overline{ba}$	yes	-

Having only bijective proofs would allow to “unfold” the induction into a bijective proof that $A_V(A)$ and $A_V(B)$ are Wilf-equivalent, for all $A \sim B$.

Bijjective proof in case (2)

$$(2) \quad a \sim b \implies PaQ \sim PbQ$$

Take $a \sim b$ and suppose that $A_V(a)$ and $A_V(b)$ are Wilf-equivalent.

Take a size-preserving bijection $\sigma : X \mapsto X^\sigma$ from $A_V(a)$ to $A_V(b)$.

Build a size-preserving bijection τ from $A_V(PaQ)$ to $A_V(PbQ)$ as follows:

Bijective proof in case (2)


$$(2) \quad a \sim b \implies PaQ \sim PbQ$$

Take $a \sim b$ and suppose that $\text{Av}(a)$ and $\text{Av}(b)$ are Wilf-equivalent.

Take a size-preserving bijection $\sigma : X \mapsto X^\sigma$ from $\text{Av}(a)$ to $\text{Av}(b)$.

Build a size-preserving bijection τ from $\text{Av}(PaQ)$ to $\text{Av}(PbQ)$ as follows:

- If X avoids PQ , then take $X^\tau = X$.
- Otherwise, apply σ to all intervals determined by the arches having one extremity between the leftmost P and the rightmost Q :



- X^τ avoids PbQ if and only if X avoids PaQ .

Analytic proof in case (4)

$$(4) \quad a \overline{bc} \sim \overline{ab} c$$

Notations: $a = \overline{A}$, $b = \overline{B}$ and $c = \overline{C}$.

F_X = the generating function of $\text{Av}(X)$.

We want that $F_{a \overline{bc}} = F_{\overline{ab} c}$.

Analytic proof in case (4)

$$(4) \quad a \overline{bc} \sim \overline{ab} c$$

Notations: $a = \overline{A}$, $b = \overline{B}$ and $c = \overline{C}$.

F_X = the generating function of $\text{Av}(X)$.

We want that $F_{a \overline{bc}} = F_{\overline{ab} c}$.

- Compute a system for $F_{a \overline{bc}}$:

$$F_{a \overline{bc}} = 1 + t F_A F_{a \overline{bc}} + t (F_{a \overline{bc}} - F_A) F_{\overline{bc}}$$

$$\text{Av}(a \overline{bc}) = \varepsilon + \underbrace{\overline{X} Y}_{X \text{ avoids } A} + \underbrace{\overline{Z} T}_{Z \text{ contains } A}$$

Analytic proof in case (4)

$$(4) \quad a \overline{bc} \sim \overline{ab} c$$

Notations: $a = \overline{A}$, $b = \overline{B}$ and $c = \overline{C}$.

F_X = the generating function of $\text{Av}(X)$.

We want that $F_{a \overline{bc}} = F_{\overline{ab} c}$.

- Compute a system for $F_{a \overline{bc}}$:

$$F_{a \overline{bc}} = 1 + tF_A F_{a \overline{bc}} + t(F_{a \overline{bc}} - F_A)F_{\overline{bc}}$$

$$F_{\overline{bc}} = 1 + tF_{bc} F_{\overline{bc}}$$

$$F_{bc} = 1 + tF_B F_{bc} + t(F_{bc} - F_B)F_c$$

$$F_c = 1 + tF_C F_c$$

Analytic proof in case (4)

$$(4) \quad a \overline{bc} \sim \overline{ab} c$$

Notations: $a = \overline{A}$, $b = \overline{B}$ and $c = \overline{C}$.

F_X = the generating function of $\text{Av}(X)$.

We want that $F_{a \overline{bc}} = F_{\overline{ab} c}$.

- Compute a system for $F_{a \overline{bc}}$:
- The solution $F_{a \overline{bc}}$ is a terrible mess depending in F_A , F_B and F_C

Analytic proof in case (4)

$$(4) \quad a\overline{bc} \sim \overline{ab}c$$

Notations: $a = \overline{A}$, $b = \overline{B}$ and $c = \overline{C}$.

F_X = the generating function of $\text{Av}(X)$.

We want that $F_{a\overline{bc}} = F_{\overline{ab}c}$.

- Compute a system for $F_{a\overline{bc}}$:
- The solution $F_{a\overline{bc}}$ is a terrible mess depending in F_A , F_B and F_C ... but symmetric in F_A , F_B and F_C !
- Consequently, $F_{a\overline{bc}} = F_{c\overline{ab}} = F_{\overline{ab}c}$.

Analytic proof in case (4)

$$(4) \quad a \overline{bc} \sim \overline{ab} c$$

Notations: $a = \overline{A}$, $b = \overline{B}$ and $c = \overline{C}$.

F_X = the generating function of $\text{Av}(X)$.

We want that $F_{a \overline{bc}} = F_{\overline{ab} c}$.

- Compute a system for $F_{a \overline{bc}}$:
- The solution $F_{a \overline{bc}}$ is a terrible mess depending in F_A , F_B and F_C ... but symmetric in F_A , F_B and F_C !
- Consequently, $F_{a \overline{bc}} = F_{c \overline{ab}} = F_{\overline{ab} c}$.
- Using $F_{\overline{X}} = 1/(1 - tF_X)$, we can write:

$$F_{a \overline{bc}} = \frac{1 - t(F_a F_b + F_b F_c + F_c F_a - F_a F_b F_c)}{1 - t(F_a + F_b + F_c - F_a F_b F_c)}$$

Enumeration of cohorts

1, 1, 2, 4, 8, 16, 32, 67, 142, 307, 669, 1 478, 3 290, 7 390, 16 709...

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.
- It follows because the arch systems of π and π' are $\sim$ -equivalent, as non-plane isomorphism corresponds to rules (3) $PabQ \sim PbaQ$,
(1) $A \sim B \implies \overline{A} \sim \overline{B}$ and (2) $a \sim b \implies PaQ \sim PbQ$.

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.
- The number of non-plane forests of size $n \sim 0.440 \cdot 2.9558^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.
- The number of non-plane forests of size $n \sim 0.440 \cdot 2.9558^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Cohorts

- = equivalence classes of non-plane forests for (4) $a\overline{bc} \sim \overline{ab}c$

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.
- The number of non-plane forests of size $n \sim 0.440 \cdot 2.9558^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Cohorts

- = equivalence classes of non-plane forests for (4) $a \overline{bc} \sim \overline{ab} c$
- The generating function of cohorts is $A(t)/t$ where

$$A = t + tA + \frac{1}{t} MSet_{\geq 2}(t^2 MSet_{\geq 3}(A)) + t MSet_{\geq 3}(A)$$

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.
- The number of non-plane forests of size $n \sim 0.440 \cdot 2.9558^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Cohorts

- $=$ equivalence classes of non-plane forests for (4) $a\overline{bc} \sim \overline{ab}c$
- The number of cohorts of size $n \sim 0.455 \cdot 2.4975^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Plane forests, non-plane forests, and cohorts

Plane forests

- Bijection ψ between 231-avoiding permutations and plane forests.
- $Cat_n \sim \frac{1}{\sqrt{\pi}} \cdot 4^n \cdot n^{-3/2}$ distinct permutation classes $Av(231, \pi)$

Non-plane forests

- If $\psi(\pi)$ and $\psi(\pi')$ are isomorphic as non-plane forests, then $Av(231, \pi)$ and $Av(231, \pi')$ are Wilf-equivalent.
- The number of non-plane forests of size $n \sim 0.440 \cdot 2.9558^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Cohorts

- $=$ equivalence classes of non-plane forests for (4) $a\overline{bc} \sim \overline{ab}c$
- The number of cohorts of size $n \sim 0.455 \cdot 2.4975^n \cdot n^{-3/2}$ is an upper bound on the number of Wilf-classes of classes $Av(231, \pi)$

Moral of the story: Many Wilf-equivalences between classes $Av(231, \pi)$!

Summary of results and open questions

- $\sim$ refines Wilf-equivalence between permutation classes $\text{Av}(231, \pi)$.
- Conjecture: $\sim$ and Wilf-equivalence coincide.
- Stronger conjecture: Given two arch systems A and B both with n arches, either $A \sim B$ or $|\text{Av}_{2n-2}(A)| \neq |\text{Av}_{2n-2}(B)|$.

Summary of results and open questions

- $\sim$ refines Wilf-equivalence between permutation classes $\text{Av}(231, \pi)$.
- Conjecture: $\sim$ and Wilf-equivalence coincide.
- Stronger conjecture: Given two arch systems A and B both with n arches, either $A \sim B$ or $|\text{Av}_{2n-2}(A)| \neq |\text{Av}_{2n-2}(B)|$.
- Asymptotic enumeration of cohorts, *i.e.* equivalence classes for $\sim$.
- It is conjecturally the number of Wilf-classes of classes $\text{Av}(231, \pi)$.

Summary of results and open questions

- $\sim$ refines Wilf-equivalence between permutation classes $\text{Av}(231, \pi)$.
- Conjecture: $\sim$ and Wilf-equivalence coincide.
- Stronger conjecture: Given two arch systems A and B both with n arches, either $A \sim B$ or $|\text{Av}_{2n-2}(A)| \neq |\text{Av}_{2n-2}(B)|$.
- Asymptotic enumeration of cohorts, i.e. equivalence classes for $\sim$.
- It is conjecturally the number of Wilf-classes of classes $\text{Av}(231, \pi)$.
- Find a completely bijective proof that $\sim$ refines Wilf-equivalence.
- Improve comparison of enumerations of $\text{Av}(231, \pi)$ and $\text{Av}(231, \pi')$.

Summary of results and open questions

- $\sim$ refines Wilf-equivalence between permutation classes $\text{Av}(231, \pi)$.
- Conjecture: $\sim$ and Wilf-equivalence coincide.
- Stronger conjecture: Given two arch systems A and B both with n arches, either $A \sim B$ or $|\text{Av}_{2n-2}(A)| \neq |\text{Av}_{2n-2}(B)|$.
- Asymptotic enumeration of cohorts, i.e. equivalence classes for $\sim$.
- It is conjecturally the number of Wilf-classes of classes $\text{Av}(231, \pi)$.
- Find a completely bijective proof that $\sim$ refines Wilf-equivalence.
- Improve comparison of enumerations of $\text{Av}(231, \pi)$ and $\text{Av}(231, \pi')$.
- Conjecture: The cohort of classes $\text{Av}(231, \pi)$ that have generating function C_n (largest possible) is the cohort with the largest cardinality.

Summary of results and open questions

- $\sim$ refines Wilf-equivalence between permutation classes $\text{Av}(231, \pi)$.
- Conjecture: $\sim$ and Wilf-equivalence coincide.
- Stronger conjecture: Given two arch systems A and B both with n arches, either $A \sim B$ or $|\text{Av}_{2n-2}(A)| \neq |\text{Av}_{2n-2}(B)|$.
- Asymptotic enumeration of cohorts, i.e. equivalence classes for $\sim$.
- It is conjecturally the number of Wilf-classes of classes $\text{Av}(231, \pi)$.
- Find a completely bijective proof that $\sim$ refines Wilf-equivalence.
- Improve comparison of enumerations of $\text{Av}(231, \pi)$ and $\text{Av}(231, \pi')$.
- Conjecture: The cohort of classes $\text{Av}(231, \pi)$ that have generating function C_n (largest possible) is the cohort with the largest cardinality.
- Extension to other contexts (separable permutations, ...).